

**STUDY OF NUTRITIONAL LIMITATIONS OF WHEAT BRAN BASED CULTURE MEDIUM FOR
SCALING UP *BTK* BASED BIOPESTICIDES PRODUCTIONS.**

Rita BARSSOUM¹, Rayan NASSERDINNE¹, Gabrielle EL KASSIS¹, Jihane SAAD^{1,3},
Meriem EL GHOUL⁴, Joanna ABBOUD^{1,3}, Nancy FAYAD¹, Julien CESCUT³, César
Arturo ACEVES-LARA², Luc FILLAUDEAU², Mireille KALLASSY AWAD¹

¹ Faculty of sciences, Saint-Joseph University of Beirut, Lebanon

² Toulouse Biotechnology Institute, INSA, Toulouse, France

³ Toulouse White Biotechnology, INSA Toulouse, France

⁴ MEDIS, Nabeul, Tunisia

Bacillus thuringiensis (*Bt*), is a spore-forming Gram-positive bacterium recognized for its insecticidal activity. This latter is due to its capacity to produce crystals formed by a mix of δ - endotoxins. *Bt* is a key player in the majority of commercially available biopesticides contributing to bio-economy expansion. The main objective of the European project IPM-4-Citrus (MSCA RISE, No. 734921) is to optimize δ - endotoxins production by *Bt* in a wheat bran (WB) based culture medium through determination of bio-performances and understanding of nutritional limitations, in order to minimize the cost of biopesticide production at industrial scale. In order to understand the nutritional limitations of the WB medium during *Bt* production, WB was first sieved and classified into 4 different sizes: class 1 >850 μm , 500 μm < class 2 <850 μm , 250 μm < class 3 <500 μm and class 4 <250 μm corresponding to 1, 19.8, 51.4 and 28.2 %w/w respectively. Two *Bt* serovar *kurstaki* (*Btk*) strains (HD1 and Lip) were cultured in flask and in 2 L bioreactor. The medium was then filtered and the following tests were carried out on permeate and retentate: ashes, starch concentration (colorimetric method), total Nitrogen (Kjeldahl) and elemental composition analysis (statistical test) of the strains and of the medium, before and after inoculation. Bio performances were also examined (CFU, spores, endotoxins). WB properties before inoculation were first determined. On one hand, the moisture (~12% gwater/gdw), ash (~4.4%w/w) and the elemental composition are constant regardless of the class. On the other hand, class 4 is the richest in starch. In suspension, retentate had an initial mass balance of 80-85% that decreased after culture indicating the partial consumption of the WB insoluble fraction while permeate had a mass balance of 14-16%. that increased after culture. Fermentable fraction increased when WB particle size decreased in agreement with starch content. Starch stands as the major carbon source (0.17-0.34 g/gWB) consumed by *Btk* however hemi-cellulosic fraction was also consumed in a rate of 0.15-0.2 g/gWB. Following, we aimed to determine nutritional limitation during *Btk* culture, based on the analysis of 5 elements (C, H, O, N, S). When compared to WB, the strain's elemental composition shows a higher nitrogen content. After culture, elemental composition was analyzed in both pellet and supernatant. Nitrogen mass balance showed significant differences between permeate and retentate after culture. It decreases in pellet due to WB consumption by *Btk* and it increases in supernatant due to biomass growth and endotoxins production. These results match with data generated by Kjeldahl technic, and confirmed a threshold for accessible Nitrogen to *Btk* due to a recalcitrant hemi-cellulosic fraction. Nitrogen was identified as a limiting nutrient at flask scale. C/N ratio in permeate and retentate appears as a reliable indicator to evaluate culture progress in flask and was used to describe biokinetics in bioreactor scale.